

Pre-operative Precautions

- Consult your doctor for special instructions if you are currently taking any medications.
- Stop taking any medication unless it is absolutely necessary.
- Patients with high blood pressure or diabetes should continue their prescribed medications. However, if you are taking Aspirin or anticoagulant (blood-thinning) medications, you must discontinue them 7 days before surgery, or follow your doctor's instructions.
- Since healthy hair follicles lead to better hair restoration results, it is advisable to undergo Meso-therapy or stem cell growth factor treatments to improve scalp condition prior to surgery.
- Check with your doctor if you are using topical treatments such as Minoxidil 5% solution or anti-hair loss shampoo before surgery.
- Inform your doctor at Maxwell if you have ever had an allergic reaction to anesthesia or antibiotics.
- If you have gray or white hair, it is recommended that you dye your hair black 2-3 days before surgery, as gray and white hair are more difficult to separate during the procedure.
- Stop smoking and drinking alcohol at least 7 days before surgery.
- Do not wear makeup, jewelry, or contact lenses, and remove nail polish and acrylic nails before arriving for surgery.



This Page



WhatsApp

WhatsApp: [821048468074](https://wa.me/821048468074)

Email: consult@maxwellhairclinic.com

Address: [11F, 120 Sanggyeong Building, Teheran-ro, Gangnam gu, Seoul](#)